arXiv:1608.01992v1 [math.NT] 5 Aug 2016

2-Variable Frobenius Problem in $\mathbb{Z}[\sqrt{M}]$

Doyon Kim[‡]

Auburn University
dzk0028@auburn.edu

July 2015

Abstract

Suppose that m is a positive integer, not a perfect square. We present a formula solution to the 2-variable Frobenius problem in $\mathbb{Z}[\sqrt{m}]$ of the "first kind" ([3]).

1 Introduction

Let $\mathbb{Z}$ denote the set of integers, and $\mathbb{N}$ the set of non-negative integers; $z_1, \ldots, z_n \in \mathbb{Z}$ are relatively prime, or coprime, if they have no common divisor in $\mathbb{Z}$ other than ± 1. Given $a_1, \ldots, a_n \in \mathbb{N} \setminus \{0\}$, let $SG(a_1, \ldots, a_n) = \{\lambda_1 a_1 + \cdots + \lambda_n a_n | \lambda_1, \ldots, \lambda_n \in \mathbb{N}\}$; $SG(a_1, \ldots, a_n)$ is the *semigroup* generated by $a_1, \ldots, a_n$.

If $a_1, \ldots, a_n \in \mathbb{N} \setminus \{0\}$ are coprime, then $SG(a_1, \ldots, a_n)$ contains a tail of $\mathbb{N}$, $\{g, g+1, \ldots\} = g + \mathbb{N}$. A Frobenius problem in $\mathbb{N}$ is the following: *For coprime positive integers $a_1, \ldots, a_n$, find the smallest $g = g(a_1, \ldots, a_n)$ such*

*Key word and phrases: 2-variable Frobenius problems, linear combination, quadratic extensions.

†AMS(MOS) Subject Classification: 11D07, 11R11, 16B99

‡This work was supported by NSF grant no. 1262930, and was completed during and after the 2015 summer Research Experience for Undergraduates in Algebra and Discrete Mathematics at Auburn University.

that $g + \mathbb{N} \subseteq SG(a_1, \ldots, a_n)$. These are called Frobenius problems because Frobenius provided a beautiful proof of a formula solution in the case $n = 2$: $g(a_1, a_2) = (a_1 - 1)(a_2 - 1)$. No such formula has been found for the cases $n \geq 3$.

In [1], [2] and [3], Frobenius problems in different rings are proposed. In the original Frobenius problem in $\mathbb{Z}$, $a_1, \ldots, a_n$ are chosen from $\mathbb{N}$, and the necessary and sufficient condition of $a_1, \ldots, a_n \in \mathbb{N} \setminus \{0\}$ to have a solution of Frobenius problem, $g = g(a_1, \ldots, a_n)$ such that $g + \mathbb{N} \subseteq SG(a_1, \ldots, a_n)$, is that they are coprime. Finding such g is equivalent to describing the set $Frob(a_1, \ldots, a_n) = \{w \in \mathbb{Z} \mid w + \mathbb{N} \in SG(a_1, \ldots, a_n)\}$, because once we find g we automatically know that $Frob(a_1, \ldots, a_n) = g + \mathbb{N}$. Solving Frobenius problems in a different ring is similar to solving these problems in $\mathbb{Z}$, but there are few more things to check to ensure that the Frobenius problem has a solution.

2 Frobenius problems in $\mathbb{Z}[\sqrt{m}]$

Suppose R is a commutative ring with multiplicative identity 1. We say that a sequence $(\alpha_1, \ldots, \alpha_n) \in R^n$ spans 1 in R if and only if for some $\lambda_1, \ldots, \lambda_n \in R$, $\lambda_1\alpha_1 + \cdots + \lambda_n\alpha_n = 1$. (In other words, the ideal generated by $\alpha_1, \ldots, \alpha_n$ in R is R.)

Let m be a positive integer which is not a perfect square, and $\mathbb{Z}[\sqrt{m}] = \{a + b\sqrt{m} \mid a, b \in \mathbb{Z}\}$. Because $\sqrt{m}$ is irrational, for each $\alpha \in \mathbb{Z}[\sqrt{m}]$ there exist unique $a, b \in \mathbb{Z}$ such that $\alpha = a + b\sqrt{m}$. Let $\mathbb{Z}[\sqrt{m}]^+ = \mathbb{Z}[\sqrt{m}] \cap [0, \infty)$ and $\mathbb{N}[\sqrt{m}] = \{a + b\sqrt{m} \mid a, b \in \mathbb{N}\}$. For $\alpha_1, \ldots, \alpha_n \in \mathbb{Z}[\sqrt{m}]$, let $SG(\alpha_1, \ldots, \alpha_n) = \{\lambda_1\alpha_1 + \cdots + \lambda_n\alpha_n \mid \lambda_1, \ldots, \lambda_n \in \mathbb{N}[\sqrt{m}]\}$ and let $SG'(\alpha_1, \ldots, \alpha_n) = \{\lambda_1\alpha_1 + \cdots + \lambda_n\alpha_n \mid \lambda_1, \ldots, \lambda_n \in \mathbb{Z}[\sqrt{m}]^+\}$. Also, let $Frob(\alpha_1, \ldots, \alpha_n) = \{w \in \mathbb{Z}[\sqrt{m}] \mid w + \mathbb{N}[\sqrt{m}] \subseteq SG(\alpha_1, \ldots, \alpha_n)\}$ and let $Frob'(\alpha_1, \ldots, \alpha_n) = \{w \in \mathbb{Z}[\sqrt{m}] \mid w + \mathbb{Z}[\sqrt{m}]^+ \subseteq SG'(\alpha_1, \ldots, \alpha_n)\}$. There are two kinds of Frobenius problem in $\mathbb{Z}[\sqrt{m}]$. One kind is, for $\alpha_1, \ldots, \alpha_n \in \mathbb{N}[\sqrt{m}] \setminus \{0\}$, to describe the set $Frob(\alpha_1, \ldots, \alpha_n)$. The other kind is, for $\beta_1, \ldots, \beta_n \in \mathbb{Z}[\sqrt{m}]^+$, to describe the set $Frob'(\beta_1, \ldots, \beta_n)$.

In [3] it is shown, and is easy to see directly, that in each kind of Frobenius problem in $\mathbb{Z}[\sqrt{m}]$, for $Frob(\alpha_1, \ldots, \alpha_n) \neq \emptyset$ (or $Frob'(\alpha_1, \ldots, \alpha_n) \neq \emptyset$) it is necessary that the sequence $(\alpha_1, \ldots, \alpha_n)$ span 1 in $\mathbb{Z}[\sqrt{m}]$. In [2] it is proved that given $\beta_1, \ldots, \beta_n \in \mathbb{Z}[\sqrt{m}]^+$, $Frob'(\beta_1, \ldots, \beta_n)$ is not empty if $(\beta_1, \ldots, \beta_n)$ spans 1 in $\mathbb{Z}[\sqrt{m}]$, and in [3] it was shown that if $\beta_1, \ldots,$

$\beta_n \in \mathbb{Z}[\sqrt{m}]^+$ spans 1 then $Frob'(\beta_1, \ldots, \beta_n) = \mathbb{Z}[\sqrt{m}]^+$. So the second kind of the problem, which is on $\mathbb{Z}[\sqrt{m}]^+$, is pretty much solved.

We will focus on the first kind of Frobenius problem, with $n = 2$. Let $\alpha_1, \alpha_2 \in \mathbb{N}[\sqrt{m}] \setminus \{0\}$, and $SG(\alpha_1, \alpha_2) = \{\lambda_1\alpha_1 + \lambda_2\alpha_2 \mid \lambda_1, \lambda_2 \in \mathbb{N}[\sqrt{m}]\}$. Our goal is to describe the set $Frob(\alpha_1, \alpha_2) = \{w \in \mathbb{Z}[\sqrt{m}] \mid w + \mathbb{N}[\sqrt{m}] \subseteq SG(\alpha_1, \alpha_2)\}$.

In [2] it is proved that if $\alpha_1, \ldots, \alpha_n \in \mathbb{N}[\sqrt{m}] \setminus \{0\}$ span 1 in $\mathbb{Z}[\sqrt{m}]$, then $Frob(\alpha_1, \ldots, \alpha_n) \neq \emptyset$ if and only if some α_i has either rational or irrational part 0, and [3] presented the solutions for the cases where every $\alpha_1, \ldots, \alpha_n$ has either rational or irrational part 0. Thanks to [3], we already know that for $m \in \mathbb{N} \setminus \{n^2 | n \in \mathbb{N}\}$ and $a, b \in \mathbb{N} \setminus \{0\}$, if a and b are coprime then $Frob(a, b) = (a-1)(b-1)(1+\sqrt{m}) + \mathbb{N}[\sqrt{m}]$, and if a and bm are coprime then $Frob(a, b\sqrt{m}) = (a-1)(b\sqrt{m}-1)(1+\sqrt{m})+\mathbb{N}[\sqrt{m}]$; $Frob(a\sqrt{m}, b\sqrt{m})$ is always empty because $a\sqrt{m}, b\sqrt{m}$ cannot span 1.

So the two remaining cases are $(\alpha_1, \alpha_2) = (a, b + c\sqrt{m})$ and $(\alpha_1, \alpha_2) = (a\sqrt{m}, b+c\sqrt{m})$, where a, b and c are positive integers. For each case, we will first find the conditions for such (α_1, α_2) to span 1 and will find $Frob(\alpha_1, \alpha_2)$. This will complete the solutions of the first Frobenius problem in $\mathbb{Z}[\sqrt{m}]$ in the case $n = 2$. The answer, by the way, is that if $\alpha, \beta \in \mathbb{N}[\sqrt{m}] \setminus \{0\}$ span 1, then $Frob(\alpha, \beta) = (\alpha - 1)(\beta - 1)(\sqrt{m} + 1) + \mathbb{N}[\sqrt{m}]$, which agrees with the results in [3] in the special cases solved there.

3 $(a, b + c\sqrt{m})$

Let $a, b, c \in \mathbb{N} \setminus \{0\}$ and let $m \in \mathbb{N} \setminus \{n^2 | n \in \mathbb{N}\}$.

Lemma 1. *$(a, b+c\sqrt{m})$ spans 1 in $\mathbb{Z}[\sqrt{m}]$ if and only if $(a, b^2 - c^2m)$ spans 1 in $\mathbb{Z}$.*

Proof. Suppose $(a, b^2 - c^2m)$ spans 1 in $\mathbb{Z}$. Then

$$a\lambda + (b^2 - c^2m)\mu = 1$$

for some $\lambda, \mu \in \mathbb{Z}$. Let

$$x = \lambda + 0 \cdot \sqrt{m} = \lambda \in \mathbb{Z}[m], \quad y = (b - c\sqrt{m})\mu \in \mathbb{Z}[m].$$

Then

$$ax + (b + c\sqrt{m})y = a\lambda + (b + c\sqrt{m})(b - c\sqrt{m})\mu = a\lambda + (b^2 - c^2m)\mu = 1.$$

Now, suppose $(a, b + c\sqrt{m})$ spans 1 in $\mathbb{Z}[\sqrt{m}]$. Then

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = 1$$

for some $x + y\sqrt{m}, z + w\sqrt{m} \in \mathbb{Z}[\sqrt{m}]$. Such $x, y, z, w \in \mathbb{Z}$ satisfies

$$ax + bz + cmw = 1,$$

$$ay + cz + bw = 0.$$

Let $\gcd(a, b^2 - c^2m) = g$. We have

$$c(ax + bz + cmw) - b(ay + cz + bw) = a(cx - by) - (b^2 - c^2m)w = c$$

so $g \mid c$, and,

$$b(ax + bz + cmw) - cm(ay + cz + bw) = a(bx - cmy) + (b^2 - c^2m)z = b$$

so $g \mid b$. Since $g \mid a$, $g \mid b$ and $g \mid c$, $g \mid ax + bz + cwm = 1$. So $g = 1$, and therefore $(a, b^2 - c^2m)$ spans 1 in $\mathbb{Z}$. □

Lemma 2. *For* $a, b, c \in \mathbb{N}\backslash\{0\}$, $m \in \mathbb{N}\backslash\{n^2 | n \in \mathbb{Z}\}$ *and* $\gcd(a, b^2 - c^2m) = 1$,

$$ax + (b + c\sqrt{m})y = A + B\sqrt{m}$$

has a solution in $\mathbb{Z}[\sqrt{m}]$ *for every* $A + B\sqrt{m} \in \mathbb{Z}[\sqrt{m}]$.

Proof. Let $A + B\sqrt{m} \in \mathbb{Z}[\sqrt{m}]$. By Lemma 1,

$$ax + (b + c\sqrt{m})y = 1$$

for some $x, y \in \mathbb{Z}[\sqrt{m}]$. Then $x(A + B\sqrt{m}) \in \mathbb{Z}[\sqrt{m}]$, $y(A + B\sqrt{m}) \in \mathbb{Z}[\sqrt{m}]$ and they satisfy

$$ax(A + B\sqrt{m}) + (b + c\sqrt{m})y(A + B\sqrt{m}) = A + B\sqrt{m}.$$

□

From now on, we assume that a, b, c are positive integers, m is a positive integer that is not a perfect square, $\gcd(a, b^2 - c^2m) = 1$, and $A, B \in \mathbb{Z}$.

Lemma 3. *If* $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ *is a solution of*

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m},$$

then every other solution $(x', y', z', w') \in \mathbb{Z}^4$ *satisfies*

$$a \mid z_0 - z', \quad a \mid w_0 - w'.$$

Proof. Let $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ be a solution of

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}.$$

Then x_0, y_0, z_0, $w_0 \in \mathbb{Z}$ satisfies

$$ax_0 + bz_0 + cmw_0 = A,$$

$$ay_0 + cz_0 + bw_0 = B.$$

Let $(x', y', z', w') \in \mathbb{Z}^4$ be another solution. x', y', z', $w' \in \mathbb{Z}$ also satisfies

$$ax' + bz' + cmw' = A,$$

$$ay' + cz' + bw' = B.$$

So we have

$$a(x_0 - x') + b(z_0 - z') + cm(w_0 - w') = 0,$$
$$a(y_0 - y') + c(z_0 - z') + b(w_0 - w') = 0.$$

Since

$$ab(x_0 - x') = -b^2(z_0 - z') - bcm(w_0 - w')$$

and

$$acm(y_0 - y') = -c^2m(z_0 - z') - bcm(w_0 - w'),$$

we have

$$a(b(x_0 - x') - cm(y_0 - y')) = -(b^2 - c^2m)(z_0 - z').$$

Since $gcd(a, b^2 - c^2m) = 1$, $a \mid z_0 - z'$. Also, we have

$$ac(x_0 - x') = -bc(z_0 - z') - c^2m(w_0 - w')$$

and

$$ab(y_0 - y') = -bc(z_0 - z') - b^2(w_0 - w'),$$

so

$$a(c(x_0 - x') - b(y_0 - y')) = (b^2 - c^2m)(w_0 - w').$$

Since $gcd(a, b^2 - c^2m) = 1$, $a \mid w_0 - w'$. □

Lemma 4. *If* $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ *is a solution of*

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m},$$

then for every $k \in \mathbb{Z}$,

$$x' = x_0 - (b - cm)k, \quad y' = y_0 + (b - c)k, \quad z' = z_0 + ak, \quad w' = w_0 - ak$$

is also a solution.

Proof. The proof is straightforward. □

Lemma 5. *If* $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ *is a solution of*

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m},$$

then for every $l \in \mathbb{Z}$,

$$x' = x_0 + bl, \quad y' = y_0 + cl, \quad z' = z_0 - al, \quad w' = w_0$$

is also a solution.

Proof. The proof is straightforward. □

Corollary 1. *There is a unique solution* $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ *of*

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$$

with $0 \leq \bar{z}, \bar{w} < a$.

Proof. Let (x_0, y_0, z_0, w_0) be a solution. Let $k = \lfloor \frac{w_0}{a} \rfloor$. Then $k \in \mathbb{Z}$, and

$$x' = x_0 - (b - cm)k, \quad y' = y_0 + (b - c)k, \quad z' = z_0 + ak, \quad w' = w_0 - ak$$

is a solution. By the choice of k, $0 \leq w' < a$ and by Lemma 3, such w' is unique.
Now, let $l = \lfloor \frac{z'}{a} \rfloor$. Then $l \in \mathbb{Z}$, and

$$\bar{x} = x' + bl, \quad \bar{y} = y' + cl, \quad \bar{z} = z' - al, \quad \bar{w} = w'$$

is also a solution, and by the choice of l, $0 \leq \bar{z} < a$. By Lemma 3, such $\bar{z}$ is unique. So $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ is a solution with $0 \leq \bar{z}, \bar{w} < a$. This is a unique such solution, because if $\bar{z}$ and $\bar{w}$ are fixed, so are $\bar{x}$ and $\bar{y}$. □

Lemma 6. *$a(x+y\sqrt{m})+(b+c\sqrt{m})(z+w\sqrt{m}) = A+B\sqrt{m}$ has a solution in $\mathbb{N}[\sqrt{m}]$ if and only if the unique solution $(\bar{x},\bar{y},\bar{z},\bar{w})$ with $0 \le \bar{z},\bar{w} < a$ satisfies $0 \le \bar{x}$, $0 \le \bar{y}$.*

Proof. Suppose $\bar{x},\bar{y} \ge 0$. Then $\bar{x}+\bar{y}\sqrt{m}$, $\bar{z}+\bar{w}\sqrt{m}$ is a solution in $\mathbb{N}[\sqrt{m}]$. Now, suppose either $\bar{x} < 0$ or $\bar{y} < 0$. Let (x',y',z',w') be another solution. If $z' < \bar{z}$ then $z' < 0$, and if $w' < \bar{w}$ then $w' < 0$, by Lemma 3. If $z' \ge \bar{z}$ and $w' \ge \bar{w}$, then

$$x' = \frac{1}{a}(A - bz' - cmw') \le \frac{1}{a}(A - b\bar{z} - cm\bar{w}) = \bar{x}, \quad \text{and}$$

$$y' = \frac{1}{a}(B - cz' - bw') \le \frac{1}{a}(B - c\bar{z} - b\bar{w}) = \bar{y}$$

so either $x' < 0$ or $y' < 0$. □

Theorem 1. *If $A \ge (a-1)(b-1+cm)$ and $B \ge (a-1)(b-1+c)$, then $a(x+y\sqrt{m})+(b+c\sqrt{m})(z+w\sqrt{m}) = A+B\sqrt{m}$ has a solution in $\mathbb{N}[\sqrt{m}]$.*

Proof. Let $A \ge (a-1)(b-1+cm)$ and $B \ge (a-1)(b-1+c)$. Consider the equation $a(x+y\sqrt{m})+(b+c\sqrt{m})(z+w\sqrt{m}) = A+B\sqrt{m}$. There is a unique solution $(\bar{x},\bar{y},\bar{z},\bar{w})$ with $0 \le \bar{z},\bar{w} < a$; $(\bar{x},\bar{y},\bar{z},\bar{w})$ satisfies

$$a\bar{x} + b\bar{z} + cm\bar{w} = A,$$

$$a\bar{y} + c\bar{z} + b\bar{w} = B.$$

Since $0 \le \bar{z},\bar{w} \le a-1$, we have

$$a\bar{x} = A - b\bar{z} - cm\bar{w} \ge (a-1)(b-1+cm) - b(a-1) - cm(a-1) = 1-a$$

so

$$\bar{x} \ge -1 + \frac{1}{a} > -1 \iff \bar{x} \ge 0.$$

Likewise,

$$a\bar{y} = B - c\bar{z} - b\bar{w} \ge (a-1)(b-1+c) - c(a-1) - b(a-1) = 1-a$$

so

$$\bar{y} \ge -1 + \frac{1}{a} > -1 \iff \bar{y} \ge 0.$$

Therefore $\bar{x}+\bar{y}\sqrt{m} \in \mathbb{N}[\sqrt{m}]$, $\bar{z}+\bar{w}\sqrt{m} \in \mathbb{N}[\sqrt{m}]$ and $\bar{x}+\bar{y}\sqrt{m}$, $\bar{z}+\bar{w}\sqrt{m}$ is a solution of $a(x+y\sqrt{m})+(b+c\sqrt{m})(z+w\sqrt{m}) = A+B\sqrt{m}$ in $\mathbb{N}[\sqrt{m}]$. □

Theorem 2. $Frob(a, b + c\sqrt{m}) = (a-1)(b + c\sqrt{m} - 1)(1 + \sqrt{m}) + \mathbb{N}[\sqrt{m}]$.

Proof. Note that

$(a-1)(b+c\sqrt{m}-1)(\sqrt{m}+1) = (a-1)(b-1+cm) + (a-1)(b-1+c)\sqrt{m}$.

In view of Theorem 1, to show that

$$Frob(a, b + c\sqrt{m}) = (a-1)(b + c\sqrt{m} - 1)(\sqrt{m} + 1) + \mathbb{N}[\sqrt{m}]$$

it suffices to show that there is no $\beta \in \mathbb{Z}$ such that

$$(a-1)(b-1+cm) - 1 + \beta\sqrt{m} \in Frob(a, b + c\sqrt{m})$$

and that there is no $\alpha \in \mathbb{Z}$ such that

$$\alpha + ((a-1)(b-1+c) - 1)\sqrt{m} \in Frob(a, b + c\sqrt{m}).$$

Let

$$A = (a-1)(b-1+cm) - 1 = ab - a - b + acm - cm.$$

Take arbitrary $k \in \mathbb{N}$, and let

$$B = (a-1)(b+c) + ak.$$

Then $(-1, k, a-1, a-1)$ is the unique solution $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ of

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$$

with $0 \leq \bar{z}, \bar{w} < a$. By Lemma 6, the equation does not have a solution in $\mathbb{N}[\sqrt{m}]$. Since the choice of k was arbitrary, B can be arbitrarily large. Therefore there is no $\beta \in \mathbb{Z}$ such that

$$(a-1)(b-1+cm) - 1 + \beta\sqrt{m} \in Frob(a, b + c\sqrt{m}).$$

Now, let

$$B = (a-1)(b-1+c) - 1.$$

Take arbitrary $l \in \mathbb{N}$, and let

$$A = (a-1)(b+cm) + al.$$

Then $(l, -1, a-1, a-1)$ is the unique solution $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ of

$$a(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$$

with $0 \leq \bar{z}, \bar{w} < a$. By Lemma 6, the equation does not have a solution in $\mathbb{N}[\sqrt{m}]$. Since the choice of l was arbitrary, A can be arbitrarily large. Therefore there is no $\alpha \in \mathbb{Z}$ such that

$$\alpha + ((a-1)(b-1+c) - 1)\sqrt{m} \in Frob(a, b + c\sqrt{m}).$$

□

4 $(a\sqrt{m}, b + c\sqrt{m})$

Let $a, b, c \in \mathbb{N} \setminus \{0\}$ and let $m \in \mathbb{N} \setminus \{n^2 | n \in \mathbb{Z}\}$.

Lemma 7. *$(a\sqrt{m}, b + c\sqrt{m})$ spans 1 in $\mathbb{Z}[\sqrt{m}]$ if and only if $(am, b^2 - c^2 m)$ spans 1 in $\mathbb{Z}$.*

Proof. Suppose $(am, b^2 - c^2 m)$ spans 1 in $\mathbb{Z}$. Then

$$am\lambda + (b^2 - c^2 m)\mu = 1$$

for some $\lambda, \mu \in \mathbb{Z}$. Let

$$x = \sqrt{m}\lambda \in \mathbb{Z}[\sqrt{m}], \quad y = (b - c\sqrt{m})\mu \in \mathbb{Z}[\sqrt{m}].$$

Then

$$a\sqrt{m}x + (b + c\sqrt{m})y = 1.$$

Now, suppose $(a\sqrt{m}, b + c\sqrt{m})$ spans 1 in $\mathbb{Z}[\sqrt{m}]$. Then

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = 1$$

for some $x + y\sqrt{m}, z + w\sqrt{m} \in \mathbb{Z}[\sqrt{m}]$; then $x, y, z, w \in \mathbb{Z}$ satisfy

$$amy + bz + cmw = 1,$$

$$ax + cz + bw = 0.$$

Then $(ay + cw)m + zb = 1$ so $\gcd(m, b^2 - c^2 m) = \gcd(m, b) = 1$. Let $\gcd(a, b^2 - c^2 m) = g$. We have

$$c(amy + bz + cmw) - b(ax + cz + bw) = a(cmy - bx) - (b^2 - c^2 m)w = c$$

so $g \mid c$, and,

$$b(amy + bz + cmw) - cm(ax + cz + bw) = a(bmy - cmx) + (b^2 - c^2 m)z = b$$

so $g \mid b$. Since $g \mid a$, $g \mid b$ and $g \mid c$, $g \mid ax + bz + cwm = 1$. Therefore $g = 1$. Since $gcd(a, b^2 - c^2 m) = 1$ and $\gcd(m, b^2 - c^2 m) = 1$, we conclude that $\gcd(am, b^2 - c^2 m) = 1$. Therefore $(am, b^2 - c^2 m)$ spans 1. □

Lemma 8. *For $a, b, c \in \mathbb{N}/\{0\}$, $m \in \mathbb{N}/\{n^2 | n \in \mathbb{Z}\}$ and* $\gcd(am, b^2 - c^2 m) = 1$*,*

$$a\sqrt{m}x + (b + c\sqrt{m})y = A + B\sqrt{m}$$

has a solution in $\mathbb{Z}[\sqrt{m}]$ for every $A + B\sqrt{m} \in \mathbb{Z}[\sqrt{m}]$.

Proof. This follows directly from Lemma 7. We skip the proof, as it is almost exactly same as the proof of Lemma 2. □

From now on, we assume that a, b, c are positive integers, m is a positive integer that is not a perfect square, $\gcd(am, b^2 - c^2 m) = 1$, and $A, B \in \mathbb{Z}$.

Lemma 9. *If $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ is a solution of*

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m},$$

then every other solution $(x', y', z', w') \in \mathbb{Z}^4$ satisfies

$$am \mid z_0 - z', \quad a \mid w_0 - w'.$$

Proof. Let $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ be a solution of

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}.$$

Then x_0, y_0, z_0, w_0 satisfies

$$amy_0 + bz_0 + cmw_0 = A,$$

$$ax_0 + cz_0 + bw_0 = B.$$

Let $(x', y', z', w') \in \mathbb{Z}^4$ be another solution. x', y', z', w' satisfies

$$amy' + bz' + cmw' = A,$$

$$ax' + cz' + bw' = B.$$

So we have

$$am(y_0 - y') + b(z_0 - z') + cm(w_0 - w') = 0,$$
$$a(x_0 - x') + c(z_0 - z') + b(w_0 - w') = 0.$$

Since

$$abm(y_0 - y') = -b^2(z_0 - z') - bcm(w_0 - w')$$

and

$$acm(x_0 - x') = -c^2m(z_0 - z') - bcm(w_0 - w'),$$

we get

$$am(b(y_0 - y') - c(x_0 - x')) = -(b^2 - c^2m)(z_0 - z').$$

Since $gcd(am, b^2 - c^2m) = 1$, $am \mid z_0 - z'$. Also, we have

$$acm(y_0 - y') = -bc(z_0 - z') - c^2m(w_0 - w')$$

and

$$ab(x_0 - x') = -bc(z_0 - z') - b^2(w_0 - w'),$$

so

$$a(cm(y_0 - y') - b(x_0 - x')) = (b^2 - c^2m)(w_0 - w').$$

Since $gcd(a, b^2 - c^2m) = 1$, $a \mid w_0 - w'$. □

Lemma 10. *If* $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ *is a solution of*

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m},$$

then for every $k \in \mathbb{Z}$,

$$x' = x_0 + (b - cm)k, \quad y' = y_0 - (b - c)k, \quad z' = z_0 + amk, \quad w' = w_0 - ak$$

is also a solution.

Proof. The proof is straightforward. □

Lemma 11. *If* $(x_0, y_0, z_0, w_0) \in \mathbb{Z}^4$ *is a solution of*

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m},$$

then for every $l \in \mathbb{Z}$,

$$x' = x_0 + cml, \quad y' = y_0 + bl, \quad z' = z_0 - aml, \quad w' = w_0$$

is also a solution.

Proof. The proof is straightforward. □

Corollary 2. *There is a unique solution* $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ *of*

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$$

with $0 \leq \bar{z} < am$, $0 \leq \bar{w} < a$.

Proof. Let (x_0, y_0, z_0, w_0) be a solution. Let $k = \lfloor \frac{w_0}{a} \rfloor$. Then $k \in \mathbb{Z}$, and

$$x' = x_0 + (b - cm)k, \quad y' = y_0 - (b - c)k, \quad z' = z_0 + amk, \quad w' = w_0 - ak$$

is a solution. By the choice of k, $0 \leq w' < a$ and by Lemma 9, such w' is unique.
Now, let $l = \lfloor \frac{z'}{am} \rfloor$. Then $l \in \mathbb{Z}$, and

$$\bar{x} = x' + cml, \quad \bar{y} = y' + bl, \quad \bar{z} = z' - aml, \quad \bar{w} = w'$$

is also a solution, and by the choice of l, $0 \leq \bar{z} < am$. By Lemma 9, such $\bar{z}$ is unique. So $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ is a solution with $0 \leq \bar{z} < am$, $0 \leq \bar{w} < a$. This is a unique such solution, because if $\bar{z}$ and $\bar{w}$ are fixed, so are $\bar{x}$ and $\bar{y}$. □

Lemma 12. *$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$ has a solution in $\mathbb{N}[\sqrt{m}]$ if and only if the unique solution $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ with $0 \leq \bar{z} < am$, $0 \leq \bar{w} < a$ satisfies $0 \leq \bar{x}$, $0 \leq \bar{y}$.*

Proof. Suppose $\bar{x}, \bar{y} \geq 0$. Then $\bar{x} + \bar{y}\sqrt{m}$, $\bar{z} + \bar{w}\sqrt{m}$ is a solution in $\mathbb{N}[\sqrt{m}]$. Now, suppose either $\bar{x} < 0$ or $\bar{y} < 0$. Let (x', y', z', w') be another solution. If $z' < \bar{z}$ then $z' < 0$, and if $w' < \bar{w}$ then $w' < 0$, by Lemma 9. If $z' \geq \bar{z}$ and $w' \geq \bar{w}$, then

$$x' = \frac{1}{a}(B - cz' - bw') \leq \frac{1}{a}(B - c\bar{z} - b\bar{w}) = \bar{x}, \quad \text{and}$$

$$y' = \frac{1}{am}(A - bz' - cmw') \leq \frac{1}{am}(A - b\bar{z} - cm\bar{w}) = \bar{y}$$

so either $x' < 0$ or $y' < 0$. □

Theorem 3. *If $A \geq abm + acm - am - cm - b + 1$ and $B \geq acm + ab - a - b - c + 1$, then $a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$ has a solution in $\mathbb{N}[\sqrt{m}]$.*

Proof. Suppose that $A \geq abm + acm - am - cm - b + 1$ and $B \geq acm + ab - a - b - c + 1$. Consider the equation $a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$. There is a unique solution $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ with $0 \leq \bar{z} < am$, $0 \leq \bar{w} < a$; $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ satisfies

$$am\bar{y} + b\bar{z} + cm\bar{w} = A,$$

$$a\bar{x} + c\bar{z} + b\bar{w} = B.$$

Since $0 \le \bar{z} \le am-1$, $\bar{w} \le a-1$, we have

$$a\bar{x} = B - c\bar{z} - b\bar{w} \ge acm + ab - a - b - c + 1 - c(am-1) - b(a-1) = 1 - a$$

so

$$\bar{x} \ge -1 + \frac{1}{a} > -1 \iff \bar{x} \ge 0.$$

Likewise,

$$am\bar{y} = A - b\bar{z} - cm\bar{w} \ge abm + acm - am - cm - b + 1 - b(am-1) - cm(a-1)$$

so

$$\bar{y} \ge -1 + \frac{1}{am} > -1 \iff \bar{y} \ge 0.$$

Therefore $\bar{x} + \bar{y}\sqrt{m} \in \mathbb{N}[\sqrt{m}]$, $\bar{z} + \bar{w}\sqrt{m} \in \mathbb{N}[\sqrt{m}]$ is a solution of $a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$ in $\mathbb{N}[\sqrt{m}]$. □

Theorem 4. $Frob(a\sqrt{m}, b + c\sqrt{m}) = (a\sqrt{m} - 1)(b + c\sqrt{m} - 1)(1 + \sqrt{m}) + \mathbb{N}[\sqrt{m}]$.

Proof. Note that

$$(a\sqrt{m} - 1)(b + c\sqrt{m} - 1)(\sqrt{m} + 1)$$

$$= (abm + acm - am - cm - b + 1) + (acm + ab - a - b - c + 1)\sqrt{m}.$$

In view of Theorem 3, to show that

$$Frob(a\sqrt{m}, b + c\sqrt{m}) = (a\sqrt{m} - 1)(b + c\sqrt{m} - 1)(\sqrt{m} + 1) + \mathbb{N}[\sqrt{m}]$$

it suffices to show that there is no $\beta \in \mathbb{Z}$ such that

$$(abm + acm - am - cm - b) + \beta\sqrt{m} \in Frob(a\sqrt{m}, b + c\sqrt{m})$$

and that there is no $\alpha \in \mathbb{Z}$ such that

$$\alpha + (acm + ab - a - b - c)\sqrt{m} \in Frob(a\sqrt{m}, b + c\sqrt{m}).$$

Let

$$A = abm + acm - am - cm - b.$$

Take arbitrary $k \in \mathbb{N}$, and let

$$B = acm + ab - b - c + ak.$$

Then $(k, -1, am-1, a-1)$ is the unique solution $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ of

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$$

with $0 \leq \bar{z} < am$, $0 \leq \bar{w} < a$. By Lemma 12, the equation does not have a solution in $\mathbb{N}[\sqrt{m}]$. Since the choice of k was arbitrary, B can be arbitrarily large. Therefore there is no $\beta \in \mathbb{Z}$ such that

$$(abm + acm - am - cm - b) + \beta\sqrt{m} \in Frob(a\sqrt{m}, b + c\sqrt{m}).$$

Now, let

$$B = acm + ab - a - b - c.$$

Take arbitrary $l \in \mathbb{N}$, and let

$$A = abm + acm - cm - b + aml.$$

Then $(-1, l, am-1, a-1)$ is the unique solution $(\bar{x}, \bar{y}, \bar{z}, \bar{w})$ of

$$a\sqrt{m}(x + y\sqrt{m}) + (b + c\sqrt{m})(z + w\sqrt{m}) = A + B\sqrt{m}$$

with $0 \leq \bar{z} < am$, $0 \leq \bar{w} < a$. By Lemma 12, the equation does not have a solution in $\mathbb{N}[\sqrt{m}]$. Since the choice of l was arbitrary, A can be arbitrarily large. Therefore there is no $\alpha \in \mathbb{Z}$ such that

$$\alpha + (acm + ab - a - b - c)\sqrt{m} \in Frob(a\sqrt{m}, b + c\sqrt{m}).$$

□

This completes the solution of the Frobenius problem of the first kind in $\mathbb{Z}[\sqrt{m}]$ when $n = 2$. We now have the following corollary:

Corollary 3. *If $\alpha_1, \alpha_2 \in \mathbb{N}[\sqrt{m}] \setminus \{0\}$, α_1, α_2 span 1, and either α_i has either rational or irrational part 0, then*

$$Frob(\alpha_1, \alpha_2) = (\alpha_1 - 1)(\alpha_2 - 1)(1 + \sqrt{m}) + \mathbb{N}[\sqrt{m}].$$

Acknowledgments

Thanks are due to Kari Vaughn, who worked on this problem before the author did and achieved results by brute force that corroborate the general results here.

References

[1] Ken Dutch, Peter Johnson, Christopher Maier, and Jordan Paschke, Frobenius problems in the Gaussian integers, *Geombinatorics* 20 (January, 2011), 93-109.

[2] Peter Johnson and Nicole Looper, Frobenius problems in integral domains, *Geombinatorics* 22 (October, 2012), 71-86.

[3] Lea Beneish, Brent Holmes, Peter Johnson and Tim Lai, Two kinds of Frobenius problems in $\mathbb{Z}[\sqrt{m}]$, *International Journal of Mathematics and Computer Science* (7), no. 2, 93-100.